

The biology of dust mites and the remediation of mite allergens in allergic disease

Larry G. Arlian, PhD,^a and Thomas A. E. Platts-Mills, MD, PhD^b Dayton, Ohio, and Charlottesville, Va

In most temperate humid areas of the world, house dust mites are a major source of multiple allergens in house dust. Mite allergens sensitize and induce perennial rhinitis, asthma, or atopic dermatitis in a large portion of patients with allergic disease. There is convincing evidence that avoidance of mite allergen can effectively reduce allergic symptoms. Patients can be moved to a mite allergen-free environment, or mite and mite allergen abatement can be performed to reduce exposure in existing residences. Some knowledge of the biology of house dust mites is essential to understand the basis of the recommendations for reducing mites and mite allergens in homes and to appreciate the difficulty of eliminating house dust mites and mite allergens from homes. This article reviews key aspects of the biology of dust mites, the properties of mite allergens, recommendations for reducing mite and mite allergen concentrations in homes, and practical recommendations for treatment. (*J Allergy Clin Immunol* 2001;107:S406-13.)

Key words: House dust mite abatement, allergen remediation, mite allergy, dust mite biology

One of the most important events in the history of allergic disease was the discovery by Voorhorst et al¹ that dust mites are the major source of allergen in house dust. They went on to explain many features of the biology of mites and provided much of the theoretical argument for current strategies used for the control of dust mites. In humid areas of the world, dust mites are ubiquitous, and up to 30% of the population have positive skin test reactions to at least one dust mite species. Sensitization to allergens derived from dust mites is strongly associated with 3 diseases: asthma, perennial rhinitis, and atopic dermatitis. Evidence for the efficacy of avoidance of dust mites and their allergens in the treatment of these diseases comes from (1) experiments in which patients' symptoms improved when they were removed from their houses^{2,3} and (2) successful controlled trials of avoidance in patients' houses.⁴ These studies provide both the evidence for recommending avoidance as part of treatment and also evidence about the methods that are most effective in controlling mite allergens.

Abbreviation used
RH: Relative humidity

BIOLOGY OF HOUSE DUST MITES

Any explanation of the biology of house dust mites cannot be complete without an overview of their classification and relationship to other arthropods, a discussion of commonly found species in both the environment and in household dust, and an examination of their prevalence, life cycle, and the roles humidity and temperature play in their development.

CLASSIFICATION AND RELATIONSHIP TO OTHER ARTHROPODS

The phylum Arthropoda is comprised of 3 subphyla of living organisms: Chelicerata, Uniramia, and Crustacea. House dust mites are arthropods belonging to the subphylum Chelicerata, class Arachnida, order Acari, and suborder Astigmata (Table I). Other suborders of mites include Mesostigmata, Metastigmata (ticks), Prostigmata, and Oribatida.

The chelicerae for the house dust mite are pincer-like and consist of both movable and fixed digits. For comparison, other chelicerates have either stylet-like chelicerae for piercing (eg, spider mites and chiggers) or sickle-like chelicerae for cutting an incision in the host skin (eg, ticks).

In addition to the mites, the subphylum Chelicerata consists of other organisms, such as sea spiders, horseshoe crabs, spiders, scorpions, and harvestman spiders (daddy longlegs), whereas insects, centipedes, and millipedes all belong to the subphylum Uniramia. Therefore mites are not closely related to insects. Mite morphology and many aspects of their physiology are very different from those of insects. For example, mites excrete primarily guanine, whereas terrestrial insects excrete primarily uric acid. Because of the differences in physiology, many insecticides and growth regulators used successfully to control insect populations are ineffective for controlling dust mite populations.

MITES IN THE ENVIRONMENT

The acari are a very diverse group of organisms that have exploited some unusual microenvironments. For

From ^athe Department of Biological Sciences, Wright State University, Dayton; and ^bAsthma and Allergic Disease Center, University of Virginia, Charlottesville.

Reprint requests: Larry G. Arlian, PhD, Department of Biological Sciences, Wright State University, 235 Biological Science Bldg, Dayton, OH 45435.

Copyright © 2001 by Mosby, Inc.

0091-6749/2001 \$35.00 + 0 1/0/113670

doi:10.1067/mai.2001.113670

TABLE I. Abbreviated classification of phylum Arthropoda

Phylum Arthropoda

1. Subphylum Uniramia
 - Class Insecta (insects)
 - Class Myriapoda (centipedes, millipedes)
2. Subphylum Crustacea
 - Class Malacostraca (crayfish, lobsters, crabs)
 - Class Maxillopoda (copepods)
3. Subphylum Chelicerata (Cheliceriformes)
 - Class Merostomata (horseshoe crabs)
 - Class Pycnogonida (sea spiders)
 - Class Arachnida
 - Order Aranea (spiders)
 - Order Opiliones (daddy longlegs)
 - Order Scorpiones
 - Order Pseudoscorpiones
 - Order Solifugae (whipscorpions)
 - Order Acari
 - Suborder Mesostigmata (free-living, predaceous, and parasitic mites)
 - Suborder Prostigmata (chiggers, follicle mites)
 - Suborder Metastigmata (ticks)
 - Suborder Astigmata (house dust, storage, & scabies mites)
 - Suborder Oribatids (soil mites)

example, some species inhabit the nasal passages of birds and mammals, the trachea of insects (mites are a major pestilence of honey bees), the hair follicles of humans and other mammals, the skin of mammals (scabies mites and others), the feathers of birds, the skin scales of reptiles, and the gills of mushrooms. In addition, many species are plant parasites, some of which are major pests for citrus, apple, strawberry, bean, and wheat crops. Many species are blood- or fluid-feeding ectoparasites of birds, mammals, and reptiles (eg, chiggers and ticks) and can transmit diseases, such as scrub typhus (chiggers) and Lyme disease (ticks). Some live in soil litter, where they are predators, fungal feeders, and detritus feeders.

MITES INDUCING ALLERGIC REACTIONS

Many species of mites that humans come in contact with, besides those found in house dust, induce allergic reactions. These include the citrus red mite (*Panonychus citri*) and the 2-spotted spider mite (*Tetranychus urticae*), which are common pests in apple orchards.⁵ The mite *Hemisarcoptes cooremani*, a parasite of scale insects that is commonly found in orchards, yards, and gardens, was recently discovered to induce IgE-mediated reactions.⁶ Scabies mites (*Sarcoptes scabiei*) burrow into the skin and induce an IgE-mediated response in many infected individuals.⁷ Many species of storage mites induce IgE-mediated reactions in exposed individuals.⁸ It should come as no surprise to learn that chiggers, ticks, and other species of ectoparasitic mites of fowl, birds, mice, guinea pigs, and other mammals and predaceous mites either sensitize or induce allergic reactions in humans as well.

MITES FOUND IN HOUSEHOLD DUST (DOMESTIC MITES)

House dust mites belong to the suborder Astigmata and family Pyroglyphidae. However, mites belonging to other families are also present in house dust, and the term domestic mites includes both mites from the Pyroglyphidae family or dust mites and mites from other families. Astigmatid mites are unusual in that they lack organized respiratory systems and associated external openings for ventilation. They are aerobic and apparently exchange O₂ and CO₂ through their general body surface. They feed principally on skin scales and other organic detritus that collects in homes. The family Pyroglyphidae contains about 16 genera and 46 species.⁹⁻¹¹ A phylogenetic perspective, the evolutionary ecology, and the cladistic relationships of families and species of astigmatid mites, can be found in several reports.^{10,12-14}

Thirteen species have been found in house dust, 3 of which are very common in homes worldwide and are the major source of mite allergen. The most common of these species are *Dermatophagoides farinae*, *Dermatophagoides pteronyssinus*, and *Euroglyphus maynei*, which are found in temperate climates. In tropical climates, the storage mite *Blomia tropicalis* (Family Echmyopodidae) can be a prevalent mite in dwellings, along with other Pyroglyphid mites. In addition, other astigmatid mites (storage mites) can be found in homes and are a potent source of allergens. Most notable are species in the families Glycyphagidae (*Glycyphagus domesticus* and *Lepidoglyphus destructor*), Acaridae (*Tyrophagus putrescentiae* and *Acarus siro*), and Chortoglyphidae (*Chortoglyphus ancutatus*). Predaceous mites (eg, Cheyletus) and parasitic mites of plants (Tetranychidae [spider mites] and Tarsonemidae) can also be present in homes. The significance of these as sources of indoor allergens is yet to be determined.

PREVALENCE OF HOUSE DUST MITES

Worldwide and in the United States, the most prevalent pyroglyphid mites found in homes are *D farinae* and *D pteronyssinus*. *E maynei* may also be prevalent in some temperate geographic areas, and at times, their density may even exceed that of *D farinae* and *D pteronyssinus*. *Blomia tropicalis* is an important source of allergen in homes in the subtropics and tropics.

In a US study of homes located in 8 different geographic areas,¹⁵ most houses were coinhabited by *D farinae* and *D pteronyssinus*. However, one species usually predominated in each home and constituted greater than 70% of the total mite population. Only a few homes in each geographic area of this study had populations containing one species only. An important discovery was that within the same geographic area, the predominant species present varied between homes. These findings illustrate the need to test for both species when a patient undergoes skin testing and when using immunotherapy.

LIFE CYCLE OF HOUSE DUST MITES

House dust mites reproduce sexually. In the female the sperm is stored in a seminal vesicle and released into the oviduct to fertilize the egg during ovulation.¹⁶ The structure of the bursa copulatrix and seminal vesicle is species specific and can be used for identification of *Dermatophagoides* species.^{16,17}

The life cycle of *D farinae*, *D pteronyssinus*, and *E maynei* consists of 5 stages: egg, larva, protonymph, tritonymph, and adult. The duration of development from egg to adult and population growth is influenced by both ambient relative humidity (RH) and temperature.^{11,18-28}

For *D pteronyssinus*, laboratory studies report that the greatest population growth occurs at 25°C, with slower growth rates occurring at lower temperatures of 10°C, 15°C, and 20°C and at higher temperatures of 30°C and 35°C.²⁹⁻³¹ Arlian et al¹⁸ reported the mean \pm SD length of the life cycle to be 122.8 \pm 14.5 days at 16°C (61°F), 34.0 \pm 5.9 days at 23°C (73°F), 19.3 \pm 2.5 days at 30°C (86°F), and 15.0 \pm 2.0 days at 35°C (95°F) at 75% RH. In comparison, for *D farinae*, few mites complete the life cycle at extreme temperatures of 16°C and 35°C, but at 23°C and 30°C, lengths of the life cycle were 35.6 \pm 4.4 days and 17.5 \pm 1.2 days, respectively.²⁵

Because temperatures are not uniform in the various areas where these mites are found in homes, the rate of reproduction, development, and mite population growth vary. For example, mite populations in carpets over slab floors that remain cool develop slower than populations inhabiting mattresses or sofas. Likewise, population growth of the 2 species in specific locations may differ because of the differences in the reproductive biology of the 2 species.

Both species produce 2 to 3 eggs per day during an approximate 26- and 34-day reproductive period for *D pteronyssinus* and *D farinae*, respectively, at 23°C and 75% RH. Mean population growth for *D farinae* and *D pteronyssinus* is 17.3% and 32.5% per week, respectively, for the humidity range of 65% to 75%.³² In laboratory cultures population densities of the 2 species decline when humidity is continuously maintained at 85% RH. However, populations exposed to daily regimens of 85% RH for 2 to 8 hours alternated with regimens of 35% RH increased in inverse proportion to time in the moist air.²³ Thus although the underlying mechanism is not fully understood, this investigation suggested that too much moist air can inhibit mite growth.

IMPORTANCE OF AMBIENT RH

House dust mites are 70% to 75% water by weight, and they obtain and maintain their water balance primarily by absorbing water from the water vapor in air.³³⁻³⁶ The lowest critical humidity at which fasting *D farinae* females can gain sufficient water from the atmosphere to maintain water balance and survive is about 55% RH at 20°C to 25°C. The critical humidity is temperature dependent, and for *D farinae*, it ranges from 55% to 75% RH over the temperature range from 15°C to 35°C.^{37,38}

Dust mites obtain water from the air by secreting a hyperosmotic solution from the supracoxal glands that open just above the first pair of legs.³⁹ The secretion flows through an open trough to the preoral cavity located beneath the chelicerae, where it is ingested. The hygroscopic solution absorbs water as it flows from the gland opening to the preoral cavity.

SOURCES OF MITE ALLERGENS

Much has been learned about dust mite allergens in the last 20 years. It is clear that mite bodies and mite feces are the sources of many allergens.^{40,41} The allergens associated with mite fecal matter are enzymes that originate from the mite's digestive tract. Possible sources of other allergens include enzymes associated with the molting process that occurs as mites change from one life stage to the next. Some allergens may be components of mite saliva that is left in the environment on food substrates where mites feed. Secretions from the supracoxal glands that are involved in the active uptake of water likely contain proteins, as well as sodium and potassium chloride.³⁹ After death, soluble protein in body fluids may be released as the body disintegrates. Some proteins from all of these sources could be allergenic.

CHARACTERISTICS OF MITE ALLERGENS

Mite allergens are divided into specific groups (1-13) on the basis of their biochemical composition, sequence homology, and molecular weight. The designation for a characterized allergen is the first 3 letters of the genus, the first letter of the species name, and a number designating the order in which the allergen was isolated or the number for other already characterized allergens it matches in homology and molecular weight. The characteristics of group 1 and 2 allergens are shown in Table II.

Group 1 allergens are glycoproteins with cysteine protease activity similar to that of some plant and mammalian enzymes.^{4,42} They originate from cells lining the intestinal tract of the mite.^{43,44} Der p 1 can cleave the CD23 IgE receptor on the membrane of human B cells and the CD25 subunit of the T-cell IL-2 receptor, which enhances its allergenicity.^{45,46} Der f 1 and Der p 1 have sequence homology of 80% and have cross-reactive epitopes. However, they also have species-specific epitopes.⁴⁷ There is some evidence that the group 1 allergens may be released as preproteins⁴⁷ that are activated by glutathione found in respiratory secretions when deposited on mucus membranes.⁴²

Group 2 allergens are 14-kd nonglycosylated proteins that have high sequence homology.⁴⁸⁻⁵⁰ Der f 2 and Der p 2 show 88% sequence homology. Group 2 allergens seem to be associated with secretion from the male mite reproductive tract,⁵¹ although Lep 2 is also associated with the gut and other areas of the mite's anatomy.⁵² The mAb 1D8 that recognizes Der p 2 but not Der f 2 also binds to Eur m 2.⁵³ This indicates that Der p 2 and Eur m 2 share an epitope not present on the Der f 2 allergen.

TABLE II. Characteristics of group 1 and 2 allergens and percentage of patient reactivity to them

Allergen group	Specific allergens characterized	Molecular weight (kd)	Frequency of reactivity (%)	Homology
Group 1	Der f 1, Der p 1, Eur m 1, Der m 1	25	>90	Cysteine protease, homology similar to enzymes papain, actinidin, cathepsin H and B, bromelain, frein
Group 2	Der f 2, Der p 2, Eur m 2, Tyr p 2, Lep d 2	14	>90	Homology with primate Epididymus protein

RECOMMENDATIONS FOR REDUCING MITE AND ALLERGEN LEVELS IN HOMES

Interventions in the home should have 3 interrelated goals: (1) reduce live mite populations; (2) reduce mite allergen levels; and (3) reduce human exposure to both. The methods used in a particular home to accomplish these 3 goals depend on many factors, most notably the cost of the intervention, the ease with which it can be implemented, the importance of a particular source of exposure, the safety of any chemicals used, and the potential effectiveness (benefit) of the intervention used. The methods selected may vary depending on one or more of these factors. A comprehensive plan, especially in a household where mite populations are extremely heavy or where persons are experiencing severe allergic reactions, may use multiple methods.

Reducing indoor RH

Maintaining RH below 50% is one of the most common recommendations for reducing dust mites and their allergen levels in homes because ambient RH is the key factor that influences dust mite prevalence. Mites must obtain sufficient water from the air to survive. Laboratory studies have shown that adult mites die of dehydration in 5 to 11 days, depending on temperature (25°C-34°C), when continuously exposed to RHs of 40% or 50%.^{35,54} Field studies report that homes located in dry climates, such as those of the mountain states or upper Midwest, have few mites and little mite allergen present. Many studies^{26-28,55-58} report seasonal fluctuations in mite and allergen levels that parallel seasonal fluctuations in indoor relative humidity. In dry climates use of evaporative coolers can raise RH enough to support mite populations.^{59,60}

The use of high-efficiency dehumidifiers and air conditioners in homes has recently been shown to be both practical and effective in reducing RH and thus mite populations. It was found that in a humid temperate climate, it is possible to maintain RH below 50%, which reduces mite and allergen levels over time.⁶¹ In addition, maintaining a mean daily RH below 50%, even when RH rises above 50% for 2 to 8 hours, effectively restricts population growth of *D. farinae* and thus the production of allergen.²³ To completely prevent population growth of *D. farinae*, RH must be maintained at 35% or less for at least 22 hours per day when daily RH is 75% to 85% for the remainder of the day.²³

In temperate climates it is possible to maintain RH at less than 50% by using high-efficiency dehumidifiers and

air conditioning. In this environment this intervention may be the only control measure needed. New carpets, mattresses, pillows, and sofas should not develop mite populations in this lower humidity environment. In homes with existing mite populations, mites should die in the dry environment, and regular vacuuming and cleaning should reduce the allergen pool in these substrates over time.⁶¹ In some climates there may be major differences in mites and mite allergens between apartments and houses, simply because of the lower humidity in the apartments. Whether dehumidification is practical and effective in warm humid climates, such as that found in Florida, remains to be determined.

Using encasements

Encasing mattresses and pillows in specially manufactured protective coverings is effective in reducing exposure to house dust mites and their allergens.⁶²⁻⁶⁶ Mattress and pillow encasements are recommended for allergic patients. Encasements may be made from plastic, vapor-permeable materials, finely woven fabrics, or nonwoven synthetics.

Pore size of the fabric is important to consider when purchasing mattress or pillow encasements. Ideally, to be comfortable, these encasements should be manufactured from breathable fabrics, be vapor permeable, and be able to block both dust mites and their allergens. Dust mite allergens (Der f 1 and Der p 1) are blocked below detectable limits by fabrics with a pore size of less than 10 µm,⁶⁷ whereas the mites themselves would be blocked by pores of less than 20 µm (width of larval stage). Dust mite larvae are generally greater than 50 µm across, and therefore access of all mites will be prevented by fabrics rated as having pore sizes of 20 µm or less.

Rains et al⁶⁸ found that new synthetic pillows accumulated Der p 1 more rapidly than new feather pillows over a 1-year period of study. Surveys of allergen levels on pillows found that nonfeather pillows contained almost 5 times more allergen⁶⁹ and an 8-fold increase of Der p 1 allergen⁷⁰ in contrast to feather pillows. It may be that tightly woven fabric covers on feather pillows prohibit mite entry and colonization of the pillows. If this theory proves to be true, manufacturers should be urged to cover all pillows with tighter fabrics.

Washing, drying, and dry cleaning of bedding materials

Washing sheets, pillow cases, blankets, and mattress pads at least weekly in hot water (55°C [130°F] or higher) kills mites and removes most allergen.^{41,71,72} When

TABLE III. House dust mite allergen avoidance: Patient education

___	Assessment of home environment
___	Description of mites (including their foods)
___	Conditions for optimal growth (humidity: >55% RH; temperature: 65°F-75°F)
___	Local areas of dust mite growth (pillows, mattresses, box springs, carpet, upholstered furniture, draperies, stuffed animals/toys)
Priorities for mites/allergens avoidance	
Bedroom	
___	Encase pillows (<10 µm pore fine woven or vapor-permeable cover)
___	Encase mattress in vapor-permeable or plastic cover
___	Encase box springs in vinyl or plastic
___	Wash bedding weekly in hot (130°F) water
House	
___	Vacuum clean weekly (wear mask; leave room for 20 minutes after cleaning)
___	Ensure that vacuum cleaner has good quality bags (usually double thickness) or high-efficiency particulate air filter on air outlet
Long-term changes to decrease mites	
___	Reduce indoor relative humidity (with air conditioning, dehumidifier, or opening windows depending on climate and season)
___	Replace carpets with polished flooring (wood, vinyl, tile)
___	Replace upholstered furniture with leather, vinyl, or wood
___	Replace draperies with wipeable shades/blinds
___	Avoid living in basements

Instructions: Record date in space provided when each measure is initiated. Bring form back on next visit.

washing in hot water, precautions should be taken to prevent the possibility of scalding. Washing in warm or cold water does not kill most mites but probably removes most allergens because allergens are water soluble.^{71,73} Adding benzyl benzoate (0.03%) and eucalyptus oils (0.2%) to the wash cycle also kills mites.⁷⁴⁻⁷⁶ However, eucalyptus oils leave an odor in clothing that lasts for 2 to 3 days.

The key to killing mites in a tumble dryer is to maintain the lethal temperature for a sufficient time. Tumble drying blankets can kill all mites if a temperature greater than 55°C (130°F) is maintained for 10 minutes.⁷⁷ Chang et al⁷⁸ found that the thermal death point and time required to kill mites was about the same in air as in water and reported that mites exposed to air at 60°C died within 10 minutes.

Dry cleaning of fabrics is effective in killing mites, but it does not destroy all allergens.^{71,79}

Replacing carpets, draperies, and upholstery

Carpets, draperies, and upholstery fabrics collect detritus and hold moisture, providing an ideal habitat for mite breeding. In humid climates it is recommended that carpets be removed in favor of hard surfaces. Likewise, draperies and curtains can be replaced with blinds or shades. To further reduce infestations, fabric upholstery can be replaced with vinyl or leather covering on cushions. Wooden furniture with no fabric is also recommended.

Vacuuming carpets

In homes with carpeting where the owners are not willing to remove the carpeting or are unable to afford to do so, regular vacuuming of the carpets at least weekly and frequent replacing of the vacuum bags is recommended. Vacuum bags with 2 layers or a high-efficiency particulate air filter or central vacuum cleaners exhausted to the outside are recommended to prevent allergens from being aerosolized during the vacuuming process.^{4,80,81} Regular

vacuum cleaning removes surface mites and allergens, but it does not remove deeply imbedded allergens or reduce the number of live mites. After a mite population has been killed, regular vacuuming of carpets is required to remove mites and allergen.^{61,82} Two studies suggest that mites and allergens may be more easily removed from low pile height and less dense carpets.^{27,83}

If temperatures are sufficiently hot, steam cleaning can kill mites and remove allergens on surfaces.⁸⁴ However, steam cleaning usually does not penetrate deep enough into the carpet, into carpet padding, or into furniture fiber, where large mite populations exist and in fact may leave sufficient residual water to promote mite population growth, which produces the exact opposite effect of what is desired (Arlian LG, unpublished observations).^{84,85}

Freezing soft toys and small items

Freezing (−17°C to −20°C) soft toys and small items (pillows and special clothing) for at least 24 hours is an effective method of killing mite populations on these objects (Arlian LG, unpublished observations).⁸⁷ After freezing in a domestic freezer, these items can then be washed to remove the dead mites and allergens. Leaving mattresses and pillows outside for at least 24 hours is a recommended method for killing mites in cold climates.

Air cleaning/filtration

Mite allergens are mainly associated with dust particles that are greater than 20 µm in diameter. These particles may become airborne as a result of disturbances, but they settle quickly.^{41,87-89} Therefore air cleaning or filtration in undisturbed spaces probably captures little mite allergen and should not be generally recommended.

Duct cleaning

No extensive studies have been performed to justify the use of duct cleaning as a remediation for household

dust mites and their allergens. This procedure is likely unnecessary to remove dust mites or mite allergen because mites do not normally reside in HVAC systems. A recently reported preliminary study found little allergen in dust collected from intake and return vent filters.⁹⁰ However, thorough studies are still needed to determine whether duct cleaning is beneficial for reducing mite allergen in the air blown from heating vents. Duct cleaning may be beneficial for removing dust, debris, and possibly mold and mold allergens from HVAC systems.

Using ozone generators

Studies are also needed to determine the efficiency and safety of ozone generators in killing mites or denaturing mite allergen. At present, there have been no published studies to show that ozone can influence mite allergen.

Using chemicals

Controlled studies have given mixed results on the use of chemicals in controlling dust mites and their allergen. Some studies show a reduction in allergen concentrations,⁹¹⁻⁹³ and others show insufficient reduction.^{81,94-96} The key considerations in the use of chemicals indoors relates to their safety, the efficacy of the active ingredient, and the formulation of the product. Products must be formulated properly so that they deliver active ingredients directly to where the mites live. Experiments with acaricides, such as benzyl benzoate, disodium octaborate tetrahydrate, sumethrin, and permethrin, and denaturants (tannic acid) have shown that a good active compound does not necessarily equate with good mite or allergen control.

DUST MITE AVOIDANCE AS AN INTEGRAL PART OF THE TREATMENT OF ALLERGIC DISEASE

Patients who have perennial rhinitis, asthma, or atopic dermatitis that is not controlled by simple topical or inhaled treatment and who are allergic to dust mites should be advised to control their exposure to mite allergens. Advice about exposure should be adapted according to the severity of the patient's disease, the climatic conditions of the area where the patient is living, and personal circumstances.

In most cases the advice given should include prioritization of the measures, as well as education about the problem and the objectives of the treatment (Table III). In all areas the patients need education, and it is helpful to develop some form of a checklist. The checklist will encourage patients to make decisions about which interventions they will take first, can afford, or both and also provides an incentive for them to record their progress.

Typically, all patients should be advised to cover pillows and mattresses, as well as change bedding so that it can be washed in hot water. The bedroom should be modified as much as is reasonable to decrease sites for mite growth and accumulation of allergens. Controlling humidity levels is not simple but should be discussed with patients as a medium- or long-term objective. In addition,

patients should be advised to make simple control of dust mites and other allergens a desirable feature of a new house when they are planning to move. Finally, it is important to remember that in all the successful controlled trials of mite avoidance for asthma, controlling mites in the bed and bedroom has been the primary measure.

CONCLUSION

In most temperate humid areas of the world, dust mites are the single most important source of allergens in house dust. Dust mite allergens are strongly associated with asthma, perennial rhinitis, and atopic dermatitis. Interventions to reduce mite populations and their allergens in the home can be worthwhile, especially when residents suffer from moderate-to-severe symptoms. The closest attention should be given to reducing mites and allergens where there is the greatest exposure in bedding. Other interventions should address control of mites and allergens in carpeting and furniture, but the advice needs to be tailored for individual patients.

We thank Jacqui Neal for the technical assistance in preparation of this article.

REFERENCES

1. Voorhorst R, Spieksma FTM, Varekamp N, et al. House dust mite atopy and the allergens it produces: identity with the house dust allergen. *J Allergy* 1967;39:325-9.
2. Boner AL, Niero E, Antolini I, Valletta EA, Gaburro D. Pulmonary function and bronchial reactivity in asthmatic children with house dust mite allergy during prolonged stay in the Italian Alps (Misurina 1756 m). *Ann Allergy* 1985;54:42-5.
3. Platts-Mills TAE, Tovey ER, Mitchell EB, Moszoro H, Nock P, Wilkins SR. Reduction of bronchial hyperactivity during prolonged allergen avoidance. *Lancet* 1982;2:675-8.
4. Platts-Mills TAE, Vervloet D, Thomas WR, Aalberse RC, Chapman MD. Indoor allergens and asthma: report of the third international workshop. *J Allergy Clin Immunol* 1997;100(Suppl):S1-24.
5. Kim YK, Lee MH, Jee YK, et al. Spider mite allergy in apple-cultivating farmers: European red mite (*Panonychus ulmi*) and two-spotted spider mite (*Tetranychus urticae*) may be important allergens in the development of work-related asthma and rhinitis symptoms. *J Allergy Clin Immunol* 1999;104:1285-92.
6. Arlian LG, Morgan MS, Houck MA. Allergenicity of the mite *Hemisarcoptes cooremani*. *Ann Allergy Asthma Immunol* 1999;83:529-32.
7. Arlian LG. Immunology of scabies. In: Wikel S, ed. Immunology of host-parasitic arthropods relationships. Wallingford, UK: CAB International; 1996. p. 232-58.
8. van Hage-Hamsten M, Johansson SG, Hoglund S, Tull P, Wiren A, Zetterstrom O. Storage mite allergy is common in a farming population. *Clin Allergy* 1985;15:555-64.
9. Wharton GW. Review article: house dust mites. *J Med Entomol* 1976;12:577-621.
10. O'Connor BM. Acari: astigmata. In: Parker S, ed. Synopsis and classification of living organisms. New York: McGraw-Hill; 1982. p. 46-69.
11. Hart BJ. Life cycle and reproduction of house-dust mites: environmental factors influencing mite populations. *Allergy* 1998;53:13-7.
12. Norton RA, Kethley JB, Johnston DE, O'Connor BM. Phylogenetic perspectives on genetic systems and reproductive modes of mites. In: Wrench DL, Ebbert MA, eds. Evolution and diversity of sex ratio in insects and mites. New York: Chapman & Hall; 1992. p. 8-99.
13. O'Connor BM. Evolutionary ecology of astigmatid mites. *Ann Rev Entomol* 1982;27:385-409.
14. Arlian LG. Mites are ubiquitous—are mite allergens, too? *Ann Allergy Asthma Immunol* 2000;85:161-3.

15. Arlian LG, Bernstein D, Bernstein IL, et al. Prevalence of dust mites in the homes of people with asthma living in eight different geographic areas of the United States. *J Allergy Clin Immunol* 1992;90:292-300.
16. Walz MG. Ultrastructure of the reproductive system of the house dust mites, *Dermatophagoides farinae* and *D. pteronyssinus* (Acari: Pyroglyphidae) with remarks on spermatogenesis and oogenesis. *Exp Appl Acarol* 1992;16:85-116.
17. Arlian LG. Biology and ecology of house dust mites, *Dermatophagoides* spp. and *Euroglyphus* spp. *Immunol Allergy Clin N Am* 1989;9:339-56.
18. Arlian LG, Rapp CM, Ahmed SG. Development of *Dermatophagoides pteronyssinus* (Acari: Pyroglyphidae). *J Med Entomol* 1990;27:1035-40.
19. Sakaki I, Suto C. The termination of prolonged quiescence (diapause) in the house dust mite *Dermatophagoides farinae* at room temperature. *Jpn J Sanit Zool* 1991;42:93-7.
20. Suto C, Sakaki I. Studies on the factors influencing the induction, persistence and termination of prolonged quiescence (diapause) in house dust mite, *Dermatophagoides farinae*. *Jpn J Sanit Zool* 1990;41:375-81.
21. Matsumoto K, Okamoto M, Wada Y. Effect of relative humidity on life cycle of the house dust mites, *Dermatophagoides farinae* and *D. pteronyssinus*. *Jpn J Sanit Zool* 1986;37:79-90.
22. Arlian LG, Neal JS, Vyszenski-Moher DL. Fluctuating hydrating and dehydrating relative humidities: effects on the life cycle of *Dermatophagoides farinae* (Acari: Pyroglyphidae). *J Med Entomol* 1999;36:457-61.
23. Arlian LG, Neal JS, Vyszenski-Moher DL. Reducing relative humidity to control the house dust mite *Dermatophagoides farinae*. *J Allergy Clin Immunol* 1999;104:852-6.
24. Arlian LG, Neal JS, Bacon SW. Survival, fecundity and development of *Dermatophagoides farinae* (Acari: Pyroglyphidae) at fluctuating relative humidity. *J Med Entomol* 1998;35:962-6.
25. Arlian LG, Dippold JS. Development and fecundity of *Dermatophagoides farinae* (Acari: Pyroglyphidae). *J Med Entomol* 1996;33:257-60.
26. Lang JD, Mulla MS. Seasonal dynamics of house dust mites, *Dermatophagoides* spp. in homes in southern California. *Environ Entomol* 1978;7:281-6.
27. Arlian LG, Bernstein IL, Gallagher JS. The prevalence of house dust mites, *Dermatophagoides* spp and associated environmental conditions in homes in Ohio. *J Allergy Clin Immunol* 1982;69:527-32.
28. Arlian LG, Woodford PJ, Bernstein IL, Gallagher JS. Seasonal population structure of house dust mites, *Dermatophagoides* spp. (Acari: Pyroglyphidae). *J Med Entomol* 1983;20:99-102.
29. Spieksma FTM. The house dust mite *Dermatophagoides pteronyssinus* (Trouessart, 1897). Producer of the house-dust allergen (Acari: Psoroptidae) [thesis]. Leiden, the Netherlands: Leiden State University; 1967.
30. Koekoek HHM, van Bronswijk JEMH. Temperature requirements of a house dust mite *Dermatophagoides pteronyssinus* compared with the climate in different habitats of houses. *Entomol Exp Appl* 1972;15:438-42.
31. Dobson RM. Some effects of microclimate on the longevity and development of *Dermatophagoides pteronyssinus* (Trouessart). *Acarologia* 1979;21:482-6.
32. Arlian LG, Confer PD, Rapp CM, Vyszenski-Moher DL, Chang JCS. Population dynamics of the house dust mites *Dermatophagoides farinae*, *D. pteronyssinus* and *Euroglyphus maynei* (Acari: Pyroglyphidae) at specific relative humidities. *J Med Entomol* 1998;35:46-53.
33. Arlian LG. Water balance and humidity requirements of house dust mites. *Exp Appl Acarol* 1992;6:15-35.
34. Arlian LG, Wharton GW. Kinetics of active and passive components of water exchange between air and a mite, *Dermatophagoides farinae*. *J Insect Physiol* 1974;20:1063-77.
35. Arlian LG. Dehydration and survival of the European house dust mite, *Dermatophagoides pteronyssinus*. *J Med Entomol* 1975;12:437-42.
36. Larson DG. The critical equilibrium activity of adult females of the house dust mite *Dermatophagoides farinae* Hughes [thesis]. Columbus, Ohio: The Ohio State University; 1969.
37. Arlian LG, Veselica MM. Reevaluation of the humidity requirements of the house dust mite *Dermatophagoides farinae* (Acari: Pyroglyphidae). *J Med Entomol* 1981;18:351-2.
38. Arlian LG, Veselica MM. Relationship between transpiration rate and temperature in the mite *Dermatophagoides farinae*. *Physiol Zool* 1981;55:344-54.
39. Wharton GW, Furumizo RT. Supracoxal gland secretions as a source of fresh water for Acarida. *Acarologia* 1977;19:112-16.
40. Arlian LG, Bernstein IL, Vyszenski-Moher DL, Gallagher JS. Investigations of culture medium-free house dust mites. III: Antigens and allergens of body and fecal extract of *Dermatophagoides farinae*. *J Allergy Clin Immunol* 1987;79:457-66.
41. Tovey ER, Chapman MD, Platts-Mills TAE. Mite faeces are a major source of house dust allergens. *Nature* 1981;289:592-3.
42. Stewart GA. Dust mite allergens. *Clin Rev Allergy Immunol* 1995;13:135-50.
43. Tovey ER, Baldo BA. Localization of antigens and allergens in thin sections of the house dust mite, *Dermatophagoides pteronyssinus* (Acari: Pyroglyphidae). *J Med Entomol* 1990;27:368-76.
44. Thomas B, Heap P, Carswell F. Ultrastructural localization of the allergen Der p1 in the gut of the house dust mite *Dermatophagoides pteronyssinus*. *Int Arch Allergy Appl Immunol* 1991;94:365-7.
45. Shakib F, Schulz O, Sewell H. A mite subversive: cleavage of CD23 and CD25 by Der p1 enhances allergenicity. *Immunol Today* 1998;19:313-6.
46. Schulz O, Sewell HF, Shakib F. Proteolytic cleavage of CD25, the subunit of the human T cell interleukin 2 receptor, by Der p1, a major mite allergen with cysteine protease activity. *J Exp Med* 1998;187:271-5.
47. Dilworth RJ, Chua KJ, Thomas WR. Sequence analysis of cDNA coding for a major house dust mite allergen, Der f1. *Clin Exp Allergy* 1991;21:25-32.
48. Heymann PW, Chapman MD, Aalberse RC, Fox JW, Platts-Mills TA. Antigenic and structural analysis of group II allergens (Der f2 and Der p2) from house dust mites (*Dermatophagoides* spp.). *J Allergy Clin Immunol* 1989;83:1055-67.
49. Lind P. Purification and partial characterization of two major allergens from the house dust mite *Dermatophagoides pteronyssinus*. *J Allergy Clin Immunol* 1985;76:753-61.
50. Yasueda H, Mita H, Yui Y, Shida T. Isolation and characterization of two allergens from *Dermatophagoides farinae*. *Int Arch Allergy Appl Immunol* 1986;81:214-23.
51. Thomas WR, Smith W. House dust mite allergens. *Allergy* 1998;53:821-32.
52. Varela J, Ventas P, Carreira J, Barbas JA, Gimenez-Gallego G, Polo F. Primary structure of Lep d I, the main *Lepidoglyphus destructor* allergen. *Eur J Biochem* 1994;225:93-8.
53. Morgan MS, Arlian LG, Barnes KC, Fernandez-Caldas EF. Characterization of the allergen of the house dust mite *Euroglyphus maynei*. *J Allergy Clin Immunol* 1997;100:222-8.
54. Brandt RL, Arlian LG. Mortality of house dust mites, *Dermatophagoides farinae* and *D. pteronyssinus* exposed to dehydrating conditions or selected pesticides. *J Med Entomol* 1976;13:327-52.
55. van Bronswijk JEMH, Sinha RN. Pyroglyphid mites (Acari) and house dust allergy. *J Allergy* 1971;47:31.
56. Murray AB, Zuk P. The seasonal variation in a population of house dust mites in a North American city. *J Allergy Clin Immunol* 1979;64:266-9.
57. Platts-Mills TAE, Hayden ML, Chapman MD, Wilkins SR. Seasonal variation in dust mite and grass pollen allergens in dust from the houses of patients with asthma. *J Allergy Clin Immunol* 1987;79:781-91.
58. Chew GL, Higgs KM, Gold DR, Muilenberg ML, Burge HA. Monthly measurements of indoor allergens and the influence of housing type in a northeastern US city. *Allergy* 1999;54:1058-66.
59. O'Rourke MK, Moore CL, Arlian LG. Prevalence of house dust mites from homes in the Sonoran Desert, Arizona. In: Muilenberg M, Burge H, eds. *Aerobiology*. New York: CRC Lewis Publishers; 1996. p. 67-80.
60. Ellingson AR, LeDoux RA, Vandanthan PK, Weber RW. The prevalence of *Dermatophagoides* mite allergen in Colorado homes utilizing central evaporative coolers. *J Allergy Clin Immunol* 1995;96:473-9.
61. Arlian LG, Alexander AK, Fowler BF, et al. Lowering humidity in homes reduces dust mites and their allergens. *J Allergy Clin Immunol* 107:99-104.
62. Ehnert B, Lau-Schadendorf S, Weber A, Beuttner P, Schou C, Wahn U. Reducing domestic exposure to dust mite allergen reduces bronchial hyperactivity in sensitive children with asthma. *J Allergy Clin Immunol* 1992;90:135-8.
63. Owen S, Morganstem M, Hepworth J, Woodcock A. Control of house dust mite antigen in bedding. *Lancet* 1990;335:396-7.
64. Custovic A, Simpson BM, Simpson A, Woodcock A. Low allergen environment can be achieved and maintained throughout pregnancy and in early life [abstract]. *J Allergy Clin Immunol* 1998;101:S81.
65. Nishioka K, Yasueda H, Saito H. Preventive effect of bedding encasement with microfibre fibers on mite sensitization. *J Allergy Clin Immunol* 1998;101:28-32.

66. Frederick JM, Warner JO, Jessop WJ, Enander I, Warner JA. Effect of a bed covering system in children with asthma and house dust mite hypersensitivity. *Eur Respir J* 1997;10:361-6.
67. Vaughan JW, McLaughlin TE, Perzanowski MS, Platts-Mills TAE. Evaluation of materials used for bedding encasement: effect of pore size in blocking cat and dust mite allergen. *J Allergy Clin Immunol* 1999;103:227-31.
68. Rains N, Siebers R, Crane J, Fitzharris P. House dust mites allergen (Der p1) accumulation on new synthetic and feather pillows. *Clin Exp Allergy* 1999;29:182-5.
69. Hallam C, Custovic A, Simpson B, Houghton N, Simpson A, Woodcock A. House dust mite allergens in feather and synthetic pillows. *Allergy* 1998;54(Suppl):407-8.
70. Kemp TJ, Siebers RW, Fishwick D, O'Grady GB, Fitzharris P, Crane J. House dust mite allergen in pillows. *BMJ* 1996;313:916.
71. McDonald LG, Tovey E. The role of water temperature and laundry procedures in reducing house dust mite populations and allergen content of bedding. *J Allergy Clin Immunol* 1992;90:599-608.
72. Siebers R, Pachett K, Fitzharris P, Crane J. Mite allergen (Der p1) on children's clothing. *J Allergy Clin Immunol* 1996;98:853-4.
73. Juhas KM, Arlian LG, Neal JS. Effects of laundry detergents and bleach on dust mite mortality [abstract]. *J Allergy Clin Immunol* 1998;101(Suppl):S28.
74. Bischoff ER, Fischer A, Liebenberg B, Kniest FM. Mite control with low temperature washing. I: Elimination of living mites on carpet pieces. *Clin Exp Allergy* 1996;26:945-52.
75. Bischoff ERC, Fischer A, Liebenberg B, Kniest FM. Mite control with low temperature washing. II: Elimination of living mites on clothing. *Clin Exp Allergy* 1998;28:60-5.
76. Tovey ER, McDonald LG. A simple washing procedure with eucalyptus oil for controlling house dust mites and their allergens in clothing and bedding. *J Allergy Clin Immunol* 1997;100:464-6.
77. Miller JD, Miller A. Ten minutes in a clothes dryer kills all mites in blankets [abstract]. *J Allergy Clin Immunol* 1996;97:423.
78. Chang JCS, Arlian LG, Dippold JS, Rapp CM, Vyszynski-Moher DL. Survival of the house dust mite, *Dermatophagoides farinae*, at high temperatures (40-80°C). *Indoor Air* 1998;8:34-8.
79. Vandehove T, Soler M, Birnbaum J, Charpin D, Vervloet D. Effect of dry cleaning on the mite allergen levels in blankets. *Allergy* 1993;48:264-6.
80. Vaughan JW, Woodfolk JA, Platts-Mills TA. Assessment of vacuum cleaners and vacuum cleaner bags recommended for allergic subjects. *J Allergy Clin Immunol* 1999;104:1079-83.
81. Woodfolk J, Hayden M, Miller J, Rose G, Chapman M, Platts-Mills T. Chemical treatment of carpets to reduce allergen: a detailed study of the effects of tannic acid on indoor allergens. *J Allergy Clin Immunol* 1994;94:19-26.
82. Munir AK, Einarsson R, Dreborg SK. Vacuum cleaning decreases the levels of mite allergens in house dust. *Pediatr Allergy Immunol* 1993;4:136-43.
83. Lewis RD, Breyse PN, Lees PSJ, diener-West M, Hamilton RG, Eggleston P. Factors affecting the retention of dust mite allergens on carpet. *Am Ind Hyg Assoc J* 1998;59:606-13.
84. Colloff M, Taylor C, Merrett T. The use of domestic steam cleaning for the control of house dust mites. *Clin Exp Allergy* 1995;25:1061-6.
85. Tovey ER, McDonald LG. Carpet cleaning. *Med J Aust* 1993;158:579.
86. Dodin A, Rak H. Influence of low temperature (-30°C) on the different stages of the human allergy mite *Dermatophagoides pteronyssinus* (Acari: Epidermoptidae). *J Med Entomol* 1993;30:810-1.
87. Custovic A, Simpson B, Simpson A, Hallam C, Craven M, Woodcock A. Relationship between mite, cat, and dog allergens in reservoir dust and ambient air. *Allergy* 1999;54:612-6.
88. Nelson HS, Hirsch SR, Ohman JL Jr, Platts-Mills TAE, Reed CE, Solomon WR. Recommendations for the use of residential air-cleaning devices in the treatment of respiratory diseases allergy. *J Allergy Clin Immunol* 1988;82:661-9.
89. Antonicelli L, Bilo MB, Pucci S, Schou C, Bonifazi F. Efficacy of an air-cleaning device equipped with a high efficiency particulate air filter in house dust mite respiratory allergy. *Allergy* 1991;46:594-600.
90. Tsay A, Roman MC, Chapman EL, Chapman MD. Air duct cleaning to reduce allergen levels in the home—is it worthwhile? [abstract]. *J Allergy Clin Immunol* 2000;105:S271.
91. Dietemann A, Bessot JC, Hoyer C, Ott M, Verot A, Pauli G. A double blind placebo controlled trial of solidified benzyl benzoate applied in dwellings of asthmatic patients sensitive to mites: clinical efficacy and effect on mite allergens. *J Allergy Clin Immunol* 1993;91:738-46.
92. Lau-Schadendorf S, Rusche AF, Weber AK, Buettner-Goetz P, Wahn U. Short-term effect of solidified benzyl benzoate on mite-allergen concentrations in house dust. *J Allergy Clin Immunol* 1991;87:41-7.
93. Vyszynski-Moher DL, Rapp CM, Neal JS, Martin F, Arlian LG. Management of house dust mites and their allergens by benzyl benzoate products [abstract]. *Ann Allergy Asthma Immunol* 2000;84:136.
94. Huss R, Huss M, Squire E, et al. Mite control with acaricide fails. *J Allergy Clin Immunol* 1992;94:27-32.
95. Manjra A, Berman D, Toerien A, Weinbert EG, Potter PC. The effects of a single treatment of an acaricide, Acarosan, and a detergent, Metsan, on Der p1 allergen levels in the carpets and mattresses of asthmatic children. *S Afr Med J* 1994;84:278-80.
96. Crank P, Hepworth J, Pickering C, Woodcock A. Concentrations of the domestic house dust mite allergen Der p1 after treatment with solidified benzyl benzoate (Acarosan) or liquid nitrogen. *Thorax* 1993;48:10-3.