

**Jay Lemery, MD**

From the Weill-Cornell Medical College, New York–Presbyterian Hospital, New York, NY.

0196-0644/\$-see front matter

Copyright © 2008 by the American College of Emergency Physicians.

doi:10.1016/j.annemergmed.2007.10.008

---



**Figure.** Pruritic rash on feet. Used with permission of J. Lemery, MD, Weill-Cornell Medical College, New York-Presbyterian Hospital, New York, NY.

[Ann Emerg Med. 2008;52:82.]

A 36-year-old male emergency physician presented 1 week after returning from a medical outreach trip in Guyana ([Figure](#)). Although mostly working in a clinical setting with wooden floors, he acknowledged a day playing beach volleyball with colleagues on a riverbank.

His only complaint was intense pruritus at the site of the lesions, and he was afebrile, with normal vital signs. A routine CBC count and chemistry panel results were within normal limits.

*For the diagnosis and teaching points, see page 92.  
To view the entire collection of Images in Emergency Medicine, visit [www.annemergmed.com](http://www.annemergmed.com)*

---

## IMAGES IN EMERGENCY MEDICINE

(continued from p. 82)

### DIAGNOSIS:

*Cutaneous larvae migrans.* This nematode parasitic infection, also known as creeping eruption, is caused by skin penetration and migration of nematode larvae. Often humans are infected when bare skin comes in contact with contaminated soil. The dermatologic sequelae present as intensely pruritic, serpiginous, erythematous lesions. Humans are accidental hosts, and the larvae are limited to the dermal-epidermal junction and do not penetrate the bloodstream. It is commonly seen in subtropical and tropical areas, as well as the southern United States. Diagnosis is mostly based on the classic presentation of the lesions, and skin biopsies are not always diagnostic. It is a self-limited infestation, but the intense pruritus and risk for secondary infection necessitate treatment. Thiabendazole, albendazole, or ivermectin are considered first-line anthelmintics.<sup>1,2</sup>

### REFERENCES

1. Liu L, Weller P. Trichinosis and infections with other tissue nematodes. In: Fauci A, Braunwald E, Isselbacher K, eds. Harrison's Principles of Internal Medicine. 14th ed. New York: McGraw-Hill; 1998:1207.
2. Mazzorana V, Maeyens E. Aquatic skin disorders. In: Auerbach P, ed. Wilderness Medicine. 5th ed. Philadelphia, PA: Mosby; 2007:1769.

## IMAGES IN EMERGENCY MEDICINE

(continued from p. 83)

### DIAGNOSIS:

*Eczema herpeticum.* A presumptive diagnosis of eczema herpeticum (Kaposi varicelliform eruption) was made at admission. The key features were the diffuse nature of the rash and the characteristic umbilicated lesions. The patient was admitted for intravenous acyclovir and vancomycin. Viral culture of a lesion grew herpes simplex 1. Direct fluorescent antibody of lesional fluid was positive for herpes simplex and negative for varicella. Eczema herpeticum is an uncommon complication of atopic dermatitis and other skin disorders because of impetiginization of abnormal and normal skin with herpes simplex. Mortality in the preacyclovir era has been reported as high as 9%.