



## Index

*Note:* Page numbers of article titles are in **boldface** type

### A

---

Abdominal delivery. See *Cesarean section*.

Abruptio placenta  
obstetric hemorrhage due to, 114–115

Acute normovolemic hemodilution  
before delivery, in parturients at risk of  
hemorrhage, 132–133

Adrenocorticotrophic hormone  
for postdural puncture headache, 76

Air embolism, venous  
during pregnancy, 170–173

Airway  
difficult, obstetric emergencies due to,  
119–120

American Society of Anesthesiologists (ASA)  
Closed Claims database in obstetrics,  
lessons learned from, **183–198**

Amniotic fluid embolism, 173–179

Analgesia, during labor  
combined spinal-epidural, 5–10  
continuous spinal, 4–5  
epidural, patient-controlled, 10–12  
patient-controlled  
epidural, 10–12  
inhalational, 2  
intravenous, 1–2

Anaphylactoid syndrome of pregnancy,  
173–179

Anesthesia  
ASA Closed Claims database on obstetric  
injuries due to, lessons learned  
from, **183–198**  
for patients with severe pre-eclampsia,  
156–160  
general. See *General anesthesia*.  
local. See *Local anesthesia*.  
obstetric. See *Obstetric anesthesia*.  
regional. See *Epidural anesthesia and  
Regional anesthesia*.

### Anesthesiologists

incomplete epidural block for cesarean  
section due to performance of,  
46–47

Antenatal assessment  
reducing risk of obstetric hemorrhage by,  
129–130

Antenatal care  
optimization of parturient to reduce risk  
of obstetric hemorrhage, 130–132

Anticoagulant medications  
obstetric anesthesia for patients taking,  
**99–109**

Antihypertensive agents  
in pre-eclampsia, 154–156

Aspiration, pulmonary  
of gastric contents during labor, decline  
in maternal mortality due to, 88–90

Atony, uterine  
obstetric hemorrhage due to, 112–114

Audit  
of cases of major obstetric hemorrhage,  
139–140

Autologous blood donation  
preoperative, in parturients at risk of  
hemorrhage, 131–132

### B

---

Back pain  
epidural analgesia and, **71–73**  
chemical backache, 73  
etiology, 71  
incidence, 71–73  
misconceptions about, 66

Backache. See *Back pain*.

Beta-adrenergic blocking agents  
to control hypertension in pre-eclampsia,  
154–155

- Blood patch. See *Epidural blood patch*.
- Blood salvage, intraoperative  
for hemorrhage in obstetric patients,  
134–135
- Blood transfusion. See *Transfusions, blood*.
- Breastfeeding  
epidural analgesia and, misconceptions  
about, 64–66
- Bupivacaine  
compared with new local anesthetics for  
obstetric anesthesia, **19–38**  
animal studies of, 20–25  
chemistry of, 19, 26  
clinical impact of toxicity of,  
29–30  
cost, 34  
human studies, 28–29  
motor block and, 30–31  
relative potency, 31–34  
toxicity of, 26–28
- 
- C**
- Calcium-channel blocking agents  
to control hypertension  
in pre-eclampsia, 155
- Cardiac arrest  
maternal, obstetric emergencies due  
to, 123
- Catheter defects  
incomplete epidural block for cesarean  
section due to, 44–46
- Catheter malfunction  
incomplete epidural block for cesarean  
section due to, 44–46
- Catheter migration  
incomplete epidural block for cesarean  
section due to, 44
- Catheter misplacement  
incomplete epidural block for cesarean  
section due to, 43–44
- Cauda equina syndrome  
after regional anesthesia, 82
- Caudal injections  
supplemental, after incomplete epidural  
block for cesarean section, 49–50
- Central nervous system  
changes in, in pre-eclampsia, 151–153
- Cesarean section  
anesthesia for, in patients with severe pre-  
eclampsia, 158–160  
general, 159–160  
regional, 158–159
- incomplete epidural block for, **39–57**  
causes of  
anatomic considerations, 43  
patient-related factors, 46  
technical skills or perfor-  
mance factor, 46  
technique, methodology, and  
equipment-related fac-  
tors, 43–46  
description of, 41–42  
management of  
if diagnosed preoperatively,  
47–50  
intraoperative, 50–52
- Chemical backache  
after regional anesthesia, 73
- Closed Claims database  
of obstetric anesthesia injury cases,  
lessons learned from, **183–198**
- Coagulation abnormalities  
in pre-eclampsia, 150–151
- Coagulation correction  
before delivery, in parturients at risk of  
hemorrhage, 132
- Coagulation defects  
obstetric anesthesia in patients with,  
**99–109**
- Continuous spinal analgesia  
during labor, 4–5
- Convulsions  
prevention of, in pre-eclampsia, 153
- Cost  
of ropivacaine and levobupivacaine *ver-*  
*sus* bupivacaine for obstetric anes-  
thesia, 34
- 
- D**
- Deep vein thrombosis  
during pregnancy, 166–170
- Dextran patch  
epidural, for postdural puncture head-  
ache, 77
- Drinking, during labor. See *Eating,*  
*during labor*.
- 
- E**
- Eating, during labor  
NPO policies, scientific validation of,  
**87–98**  
decline in maternal mortality from  
pulmonary aspiration, 88–90

- gastrointestinal anatomy and physiology during pregnancy, 90–91
- indirect evidence on, 93–96
- metabolic advantages of, 91–92
- outcome of labor and, 92–93
- Embolism**
- during pregnancy, **165–182**
- amniotic fluid embolism, 173–179
- venous air embolism, 170–173
- venous thromboembolism, 166–170
- Emergencies**
- obstetric, **111–125**
- fetal compromise, 117–119
- hemorrhage, 111–117
- abruptio placenta, 114–115
- genital tract lacerations, 111–112
- placenta accreta, 116
- placenta previa, 115–116
- retained placenta, 115
- uterine atony, 112–114
- uterine inversion, 117
- uterine rupture, 116–117
- special considerations, 119–123
- difficult airway, 119–120
- maternal cardiac arrest, 123
- morbid obesity, 121–123
- pregnancy-induced hypertension, 120–121
- Epidural analgesia**
- during labor
- combined spinal and, 5–10
- patient-controlled, 10–12
- for labor and vaginal delivery in patients with severe pre-eclampsia, 157–158
- misconceptions about, **59–70**
- back pain and, 66, **71–73**
- breastfeeding and, 64–66
- consequences of withholding, 66–68
- pre-eclampsia and, 63–64
- progress of labor, 60–63
- Epidural anesthesia**
- incomplete block for cesarean section, **39–57**
- causes of
- anatomic considerations, 43
- patient-related factors, 46
- technical skills or performance factor, 46
- technique, methodology, and equipment-related factors, 43–46
- description of, 41–42
- management of
- if diagnosed preoperatively, 47–50
- intraoperative, 50–52
- Epidural blood patch**
- for postdural puncture headache, 77
- dextran, 77
- prophylactic, 77
- treatment with, 77–79
- in HIV-positive patients, 79
- Erythropoietin, recombinant human**
- antenatal, in parturients at risk of hemorrhage, 130–131
- Esmolol**
- to control hypertension in pre-eclampsia, 155
- F**
- 
- Failed epidural.** See *Epidural anesthesia, incomplete block.*
- Fetal compromise**
- obstetric emergencies involving, 117–119
- choice of anesthesia, 118–119
- definition, 117–118
- in utero resuscitation, 118
- neonatal resuscitation, 119
- Fibrin glue injection**
- epidural, for postdural puncture headache, 77
- Fluid management**
- to control hypertension in pre-eclampsia, 155–156
- G**
- 
- Ganglion blocking agents**
- to control hypertension in pre-eclampsia, 154
- Gastric contents**
- pulmonary aspiration of during labor, decline in maternal mortality due to, 88–90
- Gastrointestinal anatomy**
- during pregnancy, 90–91
- General anesthesia**
- after incomplete epidural block for cesarean section, 50
- for cesarean delivery in patients with severe pre-eclampsia, 159–160
- increased use of, due to withholding of epidural analgesia during labor, 67

Genital tract lacerations  
obstetric hemorrhage due to, 112

## H

---

### Headache

postdural puncture, after regional analgesia during labor, **73–79**

Hemodilution, acute normovolemic  
before delivery, in parturients at risk of hemorrhage, 132–133

Hemodynamic profile  
in pre-eclampsia, 146–153

Hemorrhage, obstetric, 111–117  
abruptio placenta, 114–115  
genital tract lacerations, 111–112  
placenta accreta, 116  
placenta previa, 115–116  
retained placenta, 115  
steps to limit and treat, **127–144**  
antenatal assessment, 129–130  
antenatal optimization of parturient, 130–132  
at time of delivery, 133–136  
audit and review of cases, 139–140  
further treatment after stabilization, 138  
future alternatives to blood transfusion, 138–139  
immediately before delivery, 132–133  
intraoperative management, 136–138  
risks of blood transfusion, 128  
uterine atony, 112–114  
uterine inversion, 117  
uterine rupture, 116–117

Hemostasis, intravascular  
intraoperative maintenance of, in hemorrhage in obstetric patients, 136–137

Heparin, low-molecular-weight  
for venous thromboembolism during pregnancy, 169–170  
obstetric anesthesia for patients on, **99–109**

Hepatic dysfunction  
in pre-eclampsia, 149

Human immunodeficiency virus (HIV)  
epidural blood patch for postdural puncture headache in patients with, 79

Hydralazine  
to control hypertension in pre-eclampsia, 154

Hyperbaric oxygen  
for venous air embolism in pregnancy, 172–173

Hypertension  
pre-eclampsia, fluids, drugs and anesthetic management, **145–163**  
pregnancy-induced, obstetric emergencies due to, 120–121

## I

---

In utero resuscitation  
in obstetric emergencies involving fetal compromise, 118–119

Incomplete epidural block. See *Epidural anesthesia.*

Inhalational analgesia  
patient-controlled, during labor, 2

Injuries, due to obstetric anesthesia  
ASA Closed Claims database on, lessons learned from, **183–198**

Intraoperative blood salvage  
for hemorrhage in obstetric patients, 134–135

Intraoperative management  
of inadequate epidural block for cesarean section, 50–52

Intravenous analgesia  
patient-controlled, during labor, 1–2

Iron supplementation  
antenatal, in parturients at risk of hemorrhage, 130–131

## L

---

Labetalol  
to control hypertension in pre-eclampsia, 154–155

Labor. See also *Obstetric anesthesia.*  
newer techniques of analgesia during, **1–17**  
NPO during, guidelines for, **87–98**

Lacerations, genital tract  
obstetric hemorrhage due to, 112

Levobupivacaine. See also *Local anesthetics.*  
compared with bupivacaine for obstetric analgesia, **19–38**

Liability, medico-legal  
ASA Closed Claims database on obstetric anesthetic injuries, lessons learned from, **183–198**

Local anesthetics  
 new, for labor and delivery, **19–38**  
 chemistry of, 19, 26  
 compared with bupivacaine, animal studies of, 20–25  
 cost of, 34  
 human studies, 28–29  
 reduced motor block with, 30–31  
 relative potency of, 31–34  
 toxicity, 26–28

Local infiltration anesthesia  
 to supplement incomplete epidural block for cesarean section, 50

Low-molecular-weight heparin  
 for venous thromboembolism during pregnancy, 169–170  
 obstetric anesthesia for patients on, **99–109**

Lumbar sympathetic block  
 during labor, 3

## M

Magnesium sulfate  
 for prevention of convulsions in pre-eclampsia, 153

Malpractice claims  
 ASA Closed Claims database on claims involving obstetric anesthesia, **183–198**

Maternal cardiac arrest  
 obstetric emergencies due to, 123

Maternal death  
 due to pulmonary aspiration of gastric contents during labor, decline in, 88–90

Maternal obesity  
 obstetric emergencies due to, 121–123

Morbid obesity  
 maternal, obstetric emergencies due to, 121–123

Mortality, maternal  
 due to obstetric anesthesia, ASA Closed Claims database on, 184–187  
 due to pulmonary aspiration of gastric contents during labor, decline in, 88–90

Motor block, in obstetric anesthesia  
 with bupivacaine *versus* bupivacaine, 30–31

## N

Needle-induced trauma  
 nerve injury due to, after regional anesthesia, 80–81

Neonatal resuscitation  
 in obstetric emergencies involving fetal compromise, 119

Nerve injury  
 from needle, after regional anesthesia, 80–81

Neuraxial analgesia. See also *Epidural analgesia*.  
 in patients with thrombocytopenia, **99–109**  
 misconceptions about, **59–70**

Neurologic complications  
 after regional anesthesia during labor, **79–82**  
 cauda equina syndrome, 82  
 etiology of, 79–80  
 nerve injury from needle, 80–81  
 transient neurologic syndrome, 81–82

Nicardipine  
 to control hypertension in pre-eclampsia, 155

Nifedipine  
 to control hypertension in pre-eclampsia, 155

Nil per mouth (NPO)  
 during labor, guidelines for, **87–98**  
 decline in maternal mortality from pulmonary aspiration of gastric contents, 88–90  
 gastrointestinal anatomy and physiology during pregnancy, 90–91  
 indirect evidence on, 93–96  
 metabolic advantages of eating and drinking during labor, 91–92  
 outcome of labor, 92–93

Nitroglycerin  
 to control hypertension in pre-eclampsia, 154

Normovolemic hemodilution, acute  
 before delivery, in parturients at risk of hemorrhage, 132–133

NPO. See *Nil per mouth*.

**O**

## Obesity

- incomplete epidural block for cesarean section due to, 46
- maternal morbid, obstetric emergencies due to, 121–123

## Obstetric anesthesia, 1–198

- anticoagulant medications and, **99–109**
  - low-molecular-weight heparin, 105–107
  - practical recommendations, 104–105
- ASA closed claims, lessons learned from, **183–198**
- back pain after regional, **71–73**
- embolism during pregnancy, **165–182**
  - amniotic fluid embolism, 173–179
  - venous air embolism, 170–173
  - venous thromboembolism, 166–170
- headache after regional, **73–79**
- hemorrhage, steps to limit and treat, **127–144**
  - antenatal assessment, 129–130
  - antenatal optimization of parturient, 130–132
  - at time of delivery, 133–136
  - audit and review of cases, 139–140
  - further treatment after stabilization, 138
  - future alternatives to blood transfusion, 138–139
  - immediately before delivery, 132–133
  - intraoperative management, 136–138
  - risks of blood transfusion, 128
- in patients with thrombocytopenia, **99–109**
  - low-molecular-weight heparin and, 105–107
  - practical recommendations, 104–105
- in pre-eclampsia, **145–163**
  - anesthetic management, 156–160
  - epidural analgesia and, misconceptions about, 63–64
  - hemodynamic profile in, 146–153
  - treatment, 153–156
- incomplete epidural block for cesarean section, **39–57**
  - causes of
    - anatomic considerations, 43
    - patient-related factors, 46
    - technical skills or performance factor, 46
    - technique, methodology, and equipment-related factors, 43–46

- description of, 41–42
- management of
  - if diagnosed preoperatively, 47–50
  - intraoperative, 50–52
- neuraxial analgesia, misconceptions about, **59–70**
  - back pain and, 66, **71–73**
  - breastfeeding and, 64–66
  - consequences of withholding, 66–68
  - pre-eclampsia and, 63–64
  - progress of labor, 60–63
- neurologic complications after regional, **79–82**
- new local anesthetics, **19–38**
  - chemistry of, 19, 26
  - compared with bupivacaine, animal studies of, 20–25
  - cost of, 34
  - human studies, 28–29
  - reduced motor block with, 30–31
  - relative potency of, 31–34
  - toxicity, 26–28
- newer techniques of labor analgesia, **1–17**
  - patient-controlled analgesia
    - epidural, 10–12
    - inhalational, 2–3
    - intravenous, 1–2
  - regional techniques, 3–13
    - combined spinal-epidural analgesia, 5–10
    - continuous spinal analgesia, 4–5
    - epidural analgesia during labor, 10
    - lumbar sympathetic block, 3
    - paracervical block, 3–4
- NPO during labor, guidelines for, **87–98**
  - decline in maternal mortality from pulmonary aspiration of gastric contents, 88–90
  - gastrointestinal anatomy and physiology during pregnancy, 90–91
  - indirect evidence on, 93–96
  - metabolic advantages of eating and drinking during labor, 91–92
  - outcome of labor, 92–93
- obstetric emergencies and, **111–125**
  - fetal compromise, 117–119
  - hemorrhage, 111–117
    - abruptio placentae, 114–115
    - genital tract lacerations, 111–112
    - placenta accreta, 116
    - placenta previa, 115–116
    - retained placenta, 115
    - uterine atony, 112–114

- uterine inversion, 117
- uterine rupture, 116–117
- special considerations, 119–123
  - difficult airway, 119–120
  - maternal cardiac arrest, 123
  - morbid obesity, 121–123
  - pregnancy-induced hypertension, 120–121
- regional, backache, headache, and neurologic deficit after, **71–86**
- Oliguria
  - in pre-eclampsia, 149
- Oxygen, hyperbaric
  - for venous air embolism in pregnancy, 172–173
- Oxytocics
  - intraoperative, for hemorrhage in obstetric patients, 136

## P

---

- Paracervical block
  - during labor, 3–4
- Patient factors
  - incomplete epidural block for cesarean section due to, 46
- Patient-controlled analgesia, during labor
  - epidural, 10–12
  - inhalational, 2
  - intravenous, 1–2
- Placenta
  - obstetric hemorrhage involving
    - abruptio placenta, 114–115
    - placenta accreta, 116
    - placenta previa, 115–116
    - retained placenta, 115
- Placenta accreta
  - obstetric hemorrhage due to, 116
- Placenta previa
  - obstetric hemorrhage due to, 115–116
- Planning, anesthesia
  - to limit and treat hemorrhage in obstetric patients, **127–144**
- Postdural puncture headache
  - after regional analgesia during labor, **73–79**
- Posttraumatic stress disorders
  - due to withholding of epidural analgesia during labor, 67–68
- Pre-eclampsia, **145–163**
  - anesthetic management, 156–160
  - epidural analgesia and, misconceptions about, 63–64

- hemodynamic profile in, 146–153
- treatment, 153–156

- Pregnancy-induced hypertension
  - obstetric emergencies due to, 120–121
- Pregnancy. See also *Labor and Obstetric anesthesia*.
  - gastrointestinal anatomy and physiology in, 90–91
- Progress of labor
  - epidural analgesia and, misconceptions about, 60–63
- Pulmonary aspiration
  - of gastric contents during labor, decline in maternal mortality due to, 88–90
- Pulmonary embolism
  - during pregnancy, 166–170

## R

---

- Recombinant tissue plasminogen activator (r-tPA)
  - for venous thromboembolism during pregnancy, 169
- Regional analgesia
  - techniques for, during labor, 3–12
    - combined spinal-epidural analgesia, 5–10
    - continuous spinal analgesia, 4–5
    - epidural analgesia, patient-controlled, 10–12
    - lumbar sympathetic block, 3
    - paracervical block, 3–4
- Regional anesthesia
  - back pain after, **71–73**
  - headache after, **73–79**
  - neurologic complications after, **79–82**
    - cauda equina syndrome, 82
    - etiology of, 79–80
    - nerve injury from needle, 80–81
    - transient neurologic syndrome, 81–82
- Renal dysfunction
  - in pre-eclampsia, 149
- Respiratory system changes
  - in pre-eclampsia, 151
- Resuscitation
  - in obstetric emergencies involving fetal compromise
    - in utero, 118
    - neonatal, 119
- Retained placenta
  - obstetric hemorrhage due to, 114–115

Ropivacaine. See also *Local anesthetics*.  
 compared with bupivacaine for obstetric analgesia, **19–38**

## S

---

Saline injection  
 epidural, for postdural puncture headache, 77

Salvage, of blood, intraoperative  
 for hemorrhage in obstetric patients, 134–135

Single-shot spinal anesthesia  
 after incomplete epidural block for cesarean section, 47–48

Sodium nitroprusside  
 to control hypertension in pre-eclampsia, 154

Spinal anesthesia  
 after incomplete epidural block for cesarean section  
 combined spinal-epidural technique, 49  
 continuous, 49  
 single-shot, 47–48  
 combined epidural and, 5–10  
 continuous, during labor, 4–5

Spinal-epidural analgesia  
 combined, during labor, 5–10

Sumatriptan  
 for postdural puncture headache, 76

Supplemental injections  
 epidural or caudal, after incomplete epidural block for cesarean section, 49–50

## T

---

Theophylline  
 for postdural puncture headache, 76

Thrombocytopenia  
 obstetric anesthesia for patients with, **99–109**  
 low-molecular-weight heparin and, 105–107  
 practical recommendations, 104–105

Thrombolysis  
 for venous thromboembolism during pregnancy, 169–170

Tissue plasminogen activator, recombinant (r-tPA)  
 for venous thromboembolism during pregnancy, 169

Transfusions, blood  
 risks of, in obstetric cases, 128  
 future alternatives to, 138–139

Transient neurologic syndrome  
 after regional anesthesia, 81–82

Trimethaphan  
 to control hypertension in pre-eclampsia, 154

## U

---

Uterine atony  
 obstetric hemorrhage due to, 112–114

Uterine inversion  
 obstetric hemorrhage due to, 117

Uterine rupture  
 obstetric hemorrhage due to, 116–117

## V

---

Vasodilators, direct-acting  
 to control hypertension in pre-eclampsia, 154

Venous air embolism  
 during pregnancy, 170–173

Venous thromboembolism  
 during pregnancy, 166–170